

## PHYSICOCHEMICAL AND COLORIMETRIC PROPERTIES OF GREEN PEPPER (*CAPSICUM ANNUUM* L.) PUREE

Charles Tortoe\*, Papa Toah Akonor and Edna Meriku Essel  
Council for Scientific and Industrial Research-Food Research Institute,  
P. O. Box M20 Accra, Ghana  
\*Email: [ctortoe@yahoo.co.uk](mailto:ctortoe@yahoo.co.uk)

### Abstract

Pepper (*Capsicum annuum* L.) puree is an important food ingredient. The packaging material and shelf-life stability of green pepper puree was studied over a period of 27 days in three temperature treatments conditions as room temperature ( $30\pm 2^{\circ}\text{C}$ ), air condition temperature ( $20^{\circ}\text{C}$ ) and climatic chamber temperature ( $4^{\circ}\text{C}$ ). The green pepper puree was prepared in salt and oil in 1 and 5% concentrations pasteurised and stored in plastic and glass containers. The colour changes, pH, Total Soluble Solids (TSS) and Total Titratable Acids (TTA) were considered over the storage period. The addition of 1% salt or 5% oil to the puree caused a marked alteration in pH, TTA and colour parameters ( $L^* a^* b^*$ ) but not TSS. Over the storage period of 27 days, the pH and TTA differed significantly ( $p < 0.05$ ) among salt and oil purees with the latter having a higher pH and TTA. Storage condition did not influence these two indices but among packaging materials, glass showed ( $P < 0.05$ ) to better stabilize pH compared to the plastic material. TSS was fairly stable among the different packaging materials but was greatly reduced by storing at a higher temperature. At the end of the storage period, 5% oil puree was lighter in colour than 1% salt. Greenness of the purees remained stable over the period while yellowness reduced drastically by the final week of storage. Packaging and storage conditions did not influence colour parameters over the assessment period.

**Keywords:** *Capsicum annuum*, pepper puree, salt, oil, packaging material

Submitted: 11.02.2016

Reviewed: 16.05.2016

Accepted: 30.06.2016

### 1. INTRODUCTION

Pepper (*Capsicum* spp.) is often described as the "king of spices" and used as a flavour in foods. Capsicum belongs to the family *Solanaceae* (Night shade) and are mostly herbs or under herbs while some others are climbers. *Capsicum* contains about 90 genera and nearly 3,000 species (Vidyyarltie and Tripathi, 2002; Stern, 2000). There are numerous species of pepper (*Capsicum* spp.). The most common varieties are *Capsicum annuum*, *Capsicum baccatum*, *Capsicum chinense*, *Capsicum pubescens* and *Capsicum frutescens*. According to Messraen (1992), *Capsicum annuum* is not known in a wild state and species cultivated are commonly known as sweet pepper, bell pepper, cherry pepper and green pepper. In the tropics, pepper is mostly cultivated as rain fed crop but also in other regions as an irrigated crop (Maiti and Kumari, 2002). Pepper can be green, red, orange or yellow depending on the state of ripeness or level of maturity or the variety. Fully ripe peppers are slightly sweeter and have higher vitamin content than immature green pepper (Anonymous, 2009).

The most important characteristic feature of pepper is its hotness caused by the active ingredient, capsaicin, which is concentrated in the seeds and ribs of pepper (Anonymous, 1998). Pepper contains numerous chemicals that are importance for its nutritional value, taste, colour, aroma and antioxidant value. Chilli pepper contains, carbohydrate, protein, total fat and dietary fiber and no cholesterol. Among vitamins identified are vitamins A, C, E and K, thiamin, riboflavin, pyridoxine, pantothenic acid, niacin and folates. Minerals identified included calcium, copper, iron, magnesium, phosphorus and selenium (Mehmet *et al.*, 2006). Other components identified are water, carotenoids, resin, terpenoid and piperine (Yeager, 1998).

The health application of pepper is enormous. It has been used over the years as natural remedies for coughs, colds, sinusitis, bronchitis, ulcers and pain relief (Yeager, 1998). It is a major component in ointment for the relief of arthritic and neuropathic pains Nwachukwu (2007). According to Yeager (1998), capsaicin in peppers has anti-bacterial, anti-carcinogenic, analgesic and anti-diabetic

properties and help reduces low density lipoprotein (LDL) cholesterol, the type that causes diseases like stroke, high blood pressure and heart diseases.

A study by Joo *et al.* (2010) indicated that capsaicin is even capable of preventing the production of fat by stimulating production of proteins that break down fat and inhibiting the actions of those that produce fat. In the culinary sector, pepper is used in several dishes including curries, pizzas, meat and fish dishes, desserts as vital components of our diets. In salad preparations, pepper is an important vegetable (Remison, 2005; Fox *et al.*, 2005).

Paprika, which is a derivative from pepper is a mild red spice used primarily in the flavouring of garnishes, pickles, meats, barbecue sauces, ketchup, cheese, snack food, salads and sausages (Bosland, 1996). Pepper, is presently used in the production of ginger ale and ginger beer (Remison, 2005).

The postharvest handling of pepper is important to reduce its postharvest losses. Utilisation of pepper as a fresh commodity or processed requires appropriate postharvest handling to ensure quality product. Processing or preservation of pepper involves cleaning, packaging, storage and marketing of the pepper as fresh produce, canned, brined/pickled, frozen, fermented or dehydrated forms. Fresh peppers may be stored for up to 3 weeks in cool, moist conditions of about 45 to 50°F temperature and 85 to 90% relative humidity (Maiti and Kumari, 2002). Although *Capsicum annuum* is an economic value crop, its postharvest losses are high when fully ripe. Addressing this challenge, mature unripe green pepper (*Capsicum annuum*) was processed into puree in vegetable oil and sodium chloride and their physicochemical properties evaluated over a period of 27 days.

## 2. MATERIALS AND METHODS

### 2.1 Development of green pepper puree

Freshly harvested physiological matured green pepper (*Capsicum annuum* L.) was obtained from a farm in the Eastern Region of Ghana. The fresh pepper was sorted, destalked and

washed in distilled water. The cleaned green pepper was then washed in 1% sodium metabisulphite to remove any microbial load and subsequently washed in potable water to remove traces of sodium metabisulphite. The green pepper was milled into a smooth puree using a blender (Model 32BL79, Dynamics Corporation of America, New Hartford, Connecticut USA) at a ratio of 1:2 of pepper and distilled water.

### 2.2 Packaging material

The prepared puree was then pasteurized and stored in plastic and glass containers and kept at three temperature conditions of room temperature ( $30\pm 2^{\circ}\text{C}$ ), air condition temperature ( $20^{\circ}\text{C}$ ) and climatic chamber temperature ( $4^{\circ}\text{C}$ ) for a period of 27 days. The three selected temperature conditions of  $30\pm 2^{\circ}\text{C}$ ,  $20^{\circ}\text{C}$  and  $4^{\circ}\text{C}$  simulated green pepper puree stored in an environment without an air conditioner, with an air conditioner and in a refrigerator, respectively.

### 2.3 Development of green pepper puree in salt (sodium chloride) or oil

Sodium chloride (1% w:w) was added to the developed puree before pasteurization to obtain green pepper puree samples in salt. Similarly, vegetable oil (5% w:w) was added to the developed puree to obtain green pepper puree in vegetable oil for subsequent studies.

### 2.4 Physicochemical determination

Total Soluble Solids (TSS) of the purees defined as °Brix was determined using a Grant Refractometer RB62 (HANNA Instruments, Germany). The refractometer was initially calibrated with distilled water and sucrose solutions before the determinations in triplicates. The pH of the purees was measured using a 3330 Research pH Meter (Jenway Ltd., Essex, UK) by weighing 10.0g of the puree and subsequently dissolved in 100ml of distilled water. The mixture was allowed to stand for 1h and stirred before electrode was inserted for pH values. Total Titratable Acidity (TTA) was determined by titrimetry (Sadler and Murphy, 2010). An extract of the pepper puree (20ml)

was titrated against NaOH (0.1N) to a faint pink endpoint, using phenolphthalein as indicator, and the acidity expressed as citric acid (Bozkurt and Erkmen, 2004; Gabaldon-Leyva, 2007).

## 2.5 Colorimetry

Colour measurements of sample was conducted using a Minolta Chroma Meter (Model CR 310, Minolta Camera Co. Ltd. Japan), based on the  $L^* a^* b^*$  colour system in triplicates. The Chroma meter was calibrated with a standard white tile ( $L^* = 97.63$   $a^* = -0.48$   $b^* = +2.12$ ) as described in the Hunter Laboratory Manual (Hunter, 2001) before colour measurement of purees were taken.

## 2.6 Experimental design

A ratio of 1:2 freshly harvested green pepper and potable water was used to develop green pepper puree. A factorial design of  $2 \times 2 \times 3$  employed salt and oil puree, stored in plastic and glass containers and kept at room temperature ( $30 \pm 2^\circ\text{C}$ ), air condition temperature ( $20^\circ\text{C}$ ) and climatic chamber temperature ( $4^\circ\text{C}$ ) for a period of 27 days. The experiment was conducted in triplicates. Thirty-six samples were withdrawn per day for analysis. Six withdrawals were done over a period of 27 days consisting of four withdrawals in the 1<sup>st</sup> and 2<sup>nd</sup> week of storage at two days intervals. One withdrawal each was conducted on the 3<sup>rd</sup> and 4<sup>th</sup> week of storage at seven days interval. Based on the design a total of 216 samples were withdrawn for analysis during the period. The study employed green pepper puree without salt or oil as a control sample.

## 2.7 Statistical analysis

The Statistical Analysis Systems version 9.1 software packages were used to statistically analyze the data obtained for all treatments. Significance of treatment means was tested at  $p = 0.05$  probability level using Duncan's New Multiple Range Test (DNMRT).

## 3. RESULTS AND DISCUSSION

### 3.1 Physicochemical and colorimetric indexes of green pepper puree

Total soluble solids (TSS), pH, Total Titratable Acidity (TTA) and colour parameters of the green pepper purees are presented in Table 1. The pH and TSS content of the products are comparable to that obtained by Ahmed *et al.* (2002) for green chilli puree of pH 5.08 and TSS of 4.8. The TSS of control sample and the two treatments did not differ significantly ( $p > 0.05$ ). Addition of salt and oil treatments did not change the TSS of pepper puree but rather caused a change in pH, TTA and colour parameters. The treatments resulted in a decrease in pH for 1% salt and a further decrease for 5% oil (Table 1). This reduction in pH as a result of adding salt to the pepper puree is similar to earlier observations by Bozkurt and Erkmen, (2004) although different quantities of salt was reported in their study. Comparing the purees, 1% salt puree had the lowest TTA while 5% oil puree had the highest TTA. This observation shows that some organic acids from the oil contributed to the increase in TTA and reduction in pH of puree with 5% oil.

Colour is one of the most essential attributes of products. It influences products visual impression and contribute immensely to its acceptability or rejection (Maskan, 2001). The colour of the control samples and treatment samples were green, typical of matured green pepper. Green pepper contains chlorophyll and carotenoids which are responsible for its green colour (Marin *et al.*, 2004). Table 1 show that raw pepper puree, which served as control had a brighter colour compared to the purees in 1% salt and 5% oil. The puree containing 5% oil had the lowest  $L^*$  value compared to 1% salt. The addition of salt and the oil probably decreased the opacity of the puree, making it darker than the control (Mehdizadeh *et al.*, 2012; Farroni and Buera, 2011). Therefore, addition of oil decreased light penetration of the oil puree compared to the salt puree, thus samples with 5% oil had lower  $L^*$  value than the others.

The a\* and b\* colour values indicates that the colour of the purees lie within the green and yellow regions of the colour spectrum. Treatment with 1% salt was significantly ( $p < 0.05$ ) less in green than the control and 5% oil, while the 5% oil was the most yellow, compared to the other puree samples (Table 1). A decrease in green with the addition of salt has also been observed in other products Topno *et al.* (2013). However, in studies by Lau *et al.* (2000) and Ahmed *et al.* (2002) the attractive green colour of green plant products could be retained by controlling the degradation of chlorophyll into pheophytin during thermal processing. Additionally, thermal processing may lead to a considerable darkening of products. The higher yellowness of samples with 5% oil could be the result of the yellowish sheen of the vegetable oil used in the study.

### 3.2 Physicochemical and colour properties of pepper puree in storage

Physical and chemical properties of the purees during the period of storage are presented in Table 2. Within the storage period some puree samples were lost and were therefore discarded and excluded from the analysis. Generally, the pH of the pepper puree from 1% oil and 5% salt showed significant differences ( $p < 0.05$ ), with samples containing 1% salt having relatively lower pH as presented in Table 2. However, the pH of purees did not change significantly over the storage period of 4 weeks with packaging material and storage condition.

For puree samples treated with 1% salt the pH significantly ( $p < 0.05$ ) increased in the plastic packaging compared to glass. This increase in pH could be attributed to the breakdown of some organic acids and a subsequent formation of basic compounds (Bozkurt and Erkmen, 2004). This decomposition must have occurred considerably in plastic packaging than glass which is inert and impermeable to moisture and gases. The same observation, however, was not recorded for pepper puree treated with 5% oil. Storage condition did not have a significant influence on the pH of pepper puree. Generally, a steady reduction in TTA in the pepper puree samples over the 4-week storage period was realized. Purees treated with 5% oil had significantly higher acidity than 1% salt. The acidity of the oil, as a result of the production of free fatty acids and other organic acids in the course of storage probably influenced this occurrence. Packaging material and storage condition on the other hand had no significant effect on the TTA.

TSS was relatively stable over the study period, showing no remarkable change with regards to packaging material but differed significantly ( $p < 0.05$ ) in the case of storage condition and treatment for salt or oil. Storage under room temperature (R) saw the most profound decrease followed by air condition temperature (AC) and the least in climatic chamber (CC). Under room conditions, °Brix ranged between 3.0-5.0 while AC and CC ranged between 3.2-5.0 and 3.8-5.4, respectively (Table 2).

**Table 1: Physicochemical and colour indexes of green pepper puree**

Sample	Parameter					
	TSS	pH	TTA	L*	a*	b*
Control	4.0±0.0 <sup>a</sup>	5.55±0.01 <sup>a</sup>	0.073±0.001 <sup>a</sup>	51.73±1.07 <sup>a</sup>	-7.98±2.16 <sup>a</sup>	8.26±2.70 <sup>a</sup>
1% Salt	5.0±0.1 <sup>a</sup>	4.97±0.01 <sup>b</sup>	0.068±0.001 <sup>b</sup>	49.39±0.35 <sup>b</sup>	-5.69±0.16 <sup>b</sup>	8.15±0.29 <sup>a</sup>
5% Oil	4.0±0.1 <sup>a</sup>	5.33±0.01 <sup>c</sup>	0.079±0.002 <sup>c</sup>	49.08±0.14 <sup>b</sup>	-9.62±0.04 <sup>a</sup>	12.23±0.22 <sup>b</sup>

Within the same column, means bearing different superscripts are significantly different at 5% significance level

**Table 2: Quality properties of pepper puree during 4-week storage period**

Treatment			Physicochemical			Colour			
Day	Packaging	Storage Condition	TSS	pH	TTA	L*	a*	b*	
1% salt	Glass	*R	5.0±0.2	4.91±0.02	0.077±0.007	49.053±0.073	-4.548±0.061	12.108±0.110	
		CC	5.0±0.0	4.99±0.01	0.081±0.003	48.468±0.116	-6.070±0.093	14.200±0.160	
		AC	4.9±0.1	4.99±0.07	0.070±0.001	48.345±0.073	-5.533±0.099	13.925±0.112	
	3	Plastic	R	5.0±0.1	5.22±0.06	0.072±0.007	49.285±0.315	-4.433±0.252	11.550±0.189
			CC	4.9±0.1	5.07±0.07	0.087±0.006	48.315±0.253	-6.335±0.187	13.150±0.338
			AC	5.0±0.2	4.95±0.01	0.077±0.001	47.815±0.042	-5.628±0.065	12.388±0.082
	6	Glass	R	5.0±0.2	4.88±0.01	0.064±0.004	49.195±0.085	-4.328±0.109	11.923±0.254
			CC	5.0±0.2	5.02±0.01	0.072±0.001	48.508±0.205	-5.465±0.172	13.748±0.451
			AC	5.0±0.2	4.88±0.01	0.063±0.004	49.015±0.025	-5.015±0.057	13.045±0.134
		Plastic	R	4.0±0.1	5.31±0.01	0.064±0.002	50.688±0.145	-5.258±0.036	13.543±0.062
			CC	5.0±0.1	5.06±0.01	0.090±0.003	47.703±0.200	-6.365±0.196	12.818±0.311
			AC	5.0±0.2	5.03±0.01	0.070±0.003	48.100±0.124	-5.145±0.052	12.815±0.095
8	Glass	R	5.0±0.1	4.93±0.03	0.072±0.002	51.645±0.052	-3.920±0.039	10.623±0.115	
		CC	5.0±0.1	4.94±0.02	0.086±0.004	50.155±0.281	-4.878±0.117	11.370±0.050	
		AC	5.0±0.1	4.90±0.01	0.073±0.002	50.055±0.068	-4.575±0.047	11.380±0.071	
	Plastic	CC	5.0±0.1	4.99±0.01	0.082±0.001	50.738±0.238	-4.840±0.199	10.283±0.086	
		AC	5.0±0.1	4.97±0.01	0.081±0.003	50.925±0.116	-3.633±0.096	10.173±0.109	
		AC	5.0±0.1	4.97±0.01	0.081±0.003	50.925±0.116	-3.633±0.096	10.173±0.109	
15	Glass	R	3.0±0.1	5.02±0.01	0.085±0.001	50.935±0.060	-2.980±0.203	11.860±0.105	
		CC	5.1±0.1	5.10±0.01	0.078±0.003	48.393±0.023	-4.898±0.042	13.165±0.061	
27	Plastic	CC	5.1±0.2	5.10±0.10	0.128±0.007	48.778±0.061	-5.445±0.078	12.910±0.069	
		CC	5.0±0.1	4.77±0.01	0.074±0.001	53.283±0.417	-0.723±0.183	7.438±0.445	
5% oil	Glass	R	3.9±0.0 <sup>a</sup>	5.20±0.01 <sup>b</sup>	0.073±0.001 <sup>b</sup>	49.223±0.067 <sup>a</sup>	-5.513±0.036 <sup>a</sup>	14.013±0.088 <sup>a</sup>	
		CC	4.0±0.1	5.37±0.01	0.078±0.003 <sup>b</sup>	50.378±0.038	-6.163±0.053	13.678±0.047	
		AC	4.0±0.0	5.42±0.01	0.082±0.001 <sup>b</sup>	48.948±0.130	-5.603±0.126	14.078±0.176	
	3	Plastic	R	4.0±0.1	5.46±0.05	0.095±0.002 <sup>b</sup>	49.365±0.088	-6.275±0.017	13.908±0.046
			CC	4.0±0.1	5.41±0.01	0.073±0.004 <sup>b</sup>	51.333±0.099	-7.978±0.157	14.725±0.134
			AC	3.9±0.1	5.48±0.01	0.082±0.001 <sup>b</sup>	49.170±0.089	-6.488±0.078	13.708±0.121
	6	Glass	R	3.8±0.2	5.53±0.02	0.096±0.002 <sup>b</sup>	49.663±0.098	-5.550±0.060	15.585±0.159
			CC	5.4±0.3	5.39±0.01	0.077±0.001 <sup>b</sup>	50.773±0.101	-6.370±0.215	15.300±0.303
			AC	3.3±0.1	5.39±0.01	0.079±0.003 <sup>b</sup>	49.513±0.089	-5.685±0.013	15.525±0.080
	8	Plastic	R	3.4±0.1	5.72±0.01	0.099±0.003 <sup>b</sup>	48.655±0.037	-4.568±0.045	12.665±0.102
			CC	3.8±0.2	5.43±0.01	0.083±0.004 <sup>b</sup>	51.235±0.211	-7.735±0.259	14.665±0.366
			AC	3.6±0.3	5.26±0.01	0.085±0.007 <sup>b</sup>	50.783±0.148	-6.480±0.046	15.383±0.088
		Glass	R	3.0±0.3	4.83±0.01	0.129±0.004 <sup>b</sup>	54.183±0.069	-4.625±0.057	13.115±0.103
			CC	3.8±0.1	5.37±0.01	0.081±0.002 <sup>b</sup>	52.558±0.173	-5.423±0.243	12.775±0.258
			AC	3.2±0.2	5.78±0.02	0.064±0.004 <sup>b</sup>	51.568±0.086	-5.153±0.046	13.488±0.114
	15	Plastic	CC	4.0±0.1	5.39±0.01	0.092±0.001 <sup>b</sup>	52.640±0.158	-6.380±0.165	11.815±0.096
			AC	3.4±0.1	5.63±0.01	0.087±0.001 <sup>b</sup>	51.185±0.063	-5.478±0.025	13.653±0.026
			AC	3.4±0.1	5.63±0.01	0.087±0.001 <sup>b</sup>	51.185±0.063	-5.478±0.025	13.653±0.026
	27	Glass	CC	4.0±0.2	5.48±0.01	0.100±0.001 <sup>b</sup>	51.698±0.215	-6.163±0.149	14.418±0.206
			CC	4.1±0.4	5.52±0.01	0.083±0.004 <sup>b</sup>	50.368±0.060	-7.015±0.030	14.730±0.096
	27	Glass	CC	3.8±0.1	5.28±0.01	0.070±0.001 <sup>b</sup>	52.385±0.061	-3.725±0.016	10.828±0.109

\*R=Room, CC= Climatic chamber, AC= Air conditioned

Interesting, the difference between storage under AC and CC was not significant ( $p>0.05$ ). A significant reduction of °Brix in purees with 5% oil was observed over time but not for samples containing 1% salt (Table 2). However, pepper does not contain starch to sustain the synthesis of soluble sugars and

so an increase in TSS was not expected in any of the purees. The fairly steady pH and reduced acidity is indicative of the fact that fermentation did not occur in the pepper puree during storage. Colour loss or degradation is considered undesirable since it may lead to the rejection of a product. This

may result from browning reactions or breakdown of colour pigments into other forms. The behaviour of pepper puree colour in storage is also presented in Table 2. The  $L^*$  value was considerably different between 1% salt and 5% oil, with 5% oil being less coloured. However, storage condition and packaging material did not impact more colour to the pepper puree samples over the study period. Overall,  $L^*$  value showed a drift toward less colour during the third and fourth weeks of storage. Redness-greenness of the purees remained stable without marked change ( $p > 0.05$ ) observed for the different treatments, packaging material or storage condition. This suggests that there was not any significant breakdown of green colour compounds in the puree during storage. The observation also downplays the formation of brown pigments through browning reactions and ascorbic acid oxidation in purees during storage (Bozkurt and Erkmen, 2004). The yellowness or blueness, unlike red-green changed markedly over the storage period. The  $b^*$  value increased during storage (non-linear) however, its value reduced sharply in the fourth week. This reduction is indicative of loss in the yellowness of the purees in storage which is probably due to oxidation of yellow colour compounds (Ahmed, 2004). Changes that occurred in colour of the purees, nonetheless, were not influenced by packaging material or storage condition, contrary to what have been reported in other studies (Ahmed and Shivare, 2001; Sigge *et al.*, 2001; Silvica *et al.*, 2011). For retention of green colour, low temperature storage has been suggested by Sigge *et al.* (2001).

#### 4. CONCLUSIONS

Pepper purees at reduced temperature of storage in glass containers showed good shelf-stability. Although some physico-chemical properties changed over the period of storage, colour (greenness) of the purees remained relatively stable and was not influenced by packaging or storage condition.

#### 5. REFERENCES

- [1] R. D. Vidyarthi and S. C. Tripathi, A textbook of Botany. S. Chand and Company Ltd. 7361, Ram Nagar, New Delhi, 2002.
- [2] K. R. Stern, Introductory plant biology, 8th edition McGraw Hill College, New York pp630, 2000.
- [3] C. M. Messiaen, The Tropical Vegetable Garden. Macmillian (London). pp. 2, 1992.
- [4] S. Maiti and P. T. K. Kumari, Cultivation of Long Pepper, Anand Press, Anand 388001, India, 2002.
- [5] Anonymous, Power your diet. (<http://www.nutrition-and-you.com/> Power your diet), 2009.
- [6] Anonymous, Peppers: Safe Methods to Store, Preserve and Enjoy, University of California Publication 80004, USA, 1998.
- [7] O. Mehmet, B. Berna, A. Turkan and S. Zeynep Identification of pepper viruses by Das-Elisa assays in Gaziantep-Turkey. *Plant Pathology Journal*, 5:11-14, 2006.
- [8] USDA, National Nutrient Database for standard reference. <http://www.nal.usda.gov/fnic/foodcomp/search>, 2008.
- [9] S. Yeager, The Doctor's Book of Food Remedies, Rodale Inc., USA, pp. 145 – 149, 1998.
- [10] C. U. Nwachukwu, F. N. Mbagwu, A. N. Onyeji, Morphological and leaf epidermal features of *Capsicum annuum* and *Capsicum frutescens* solanaceae. *Nature and Science*, 5: 54 – 60, 2007.
- [11] J. I. Joo, D. H. Kim, J. W. Choi and J. W. Yun, Proteomic analysis for anti-obesity potential of capsaicin on white adipose tissue in rat fed with a high fat diet. *Journal of Proteome Research*, 9:2977-2987, 2010, Doi: 10.1021/pr901175w.
- [12] S. U. Remison, Arable and Vegetables Crops of the Tropics. Gift Prints Associates, Benin City. 247p, 2005.
- [13] A. J. Fox, D. D. Pozo-Insfran, J. H. Lee, S. A. Sargent and S. T. Talcott, Ripening-induced chemical and Antioxidant changes in Bell peppers as affected by harvest maturity and postharvest ethylene exposure. *HortScience*, 40:732-736, 2005.
- [14] P. W. Bosland. Capsicums: Innovative uses of an ancient crop. In: Janick, J. (ed). Progress in new Crops. ASHS Press, Arlington, Va. pp479 – 487, 1996.
- [15] G. D. Sadler and P.A. Murphy, pH and Titratable Acidity. In: Nielsen, S (ed), Food Analysis 4<sup>th</sup> Edition, Springer Science and Business Media, LLC, New York. pp. 219 – 238, 2010.
- [16] H. Bozkurt and O. Erkmen, Effects of production techniques on the quality of hot pepper paste. *Journal of Food Engineering* 64,173-178, 2004.
- [17] C. A. Gabaldon-Leyva, A. Quintero-Ramos, J. Barnard, R. R. Balandran-Quintana, R. Talamas-Abbud, and J. Jimenez-Castro, Effect of ultrasound on the mass transfer and physical changes in brine bell

- pepper at different temperatures. *Journal of Food Engineering*, 81:374-379, 2007.
- [18] Hunter Laboratory Manual Hunter, Associate Laboratory Universal Software. Version 3.8. ISO 9001 certified. Reston, VA, USA, 2001.
- [19] J. Ahmed, U. S. Shivhare and S. Debnath, Colour degradation and rheology of green chilli puree during thermal processing. *International Journal of Food Science and Technology*, 37:57-63, 2002.
- [20] M. Maskan, Kinetics of colour change of kiwifruits during hot air and microwave drying. *Journal of Food Engineering*, 48:169-175, 2001.
- [21] A. Marin, F. Ferreres, F. A. Thomas-Barberan, M. I. Gil. Characterization and quantitation of antioxidant constituents of sweet pepper (*Capsicum annum* L). *Journal of Agriculture and Food Chemistry*, 52:3861-3869, 2004.
- [22] T. Mehdizadeh, H. Tajik, S. M. R. Rohani and A. R. Oromiehie, Antibacterial, antioxidant and optical properties of edible starch-chitosan composite film containing *Thymus kotschyianus* essential oil. *Veterinary Research Forum*, 3:167-173, 2012.
- [23] A. Farroni and M. D. P. Buera, Water, Maillard reaction and food appearance. The world of food science. <http://www.worldfoodscience.org/cms/?pid=1005741>: accessed 16<sup>th</sup> August 2013, 2011.
- [24] P. N. Topno, V. Vinothini, S. H. Jayaprakash, V. Varadaiah, S. H. Sheshagiri, Srinivas, P. M. and Naidu, M. M. Ginger-garlic paste in retort pouches and its quality. *Journal of Food Process Engineering*, 36:1-8. 2013.
- [25] Lau, M. H., Tang, J. and Swanson, B. G. Kinetics of textural and colour changes in green asparagus during thermal treatments. *Journal of Food Engineering*, 45:231-236, 2000.
- [26] J. Ahmed, Rheological behavior and colour changes in ginger paste during storage. *International Journal of Food Science and Technology*, 39:325-330, 2004.
- [27] J. Ahmed and U. S Shivare, Thermal kinetics of colour change, rheology and storage characteristics of Garlic puree/paste. *Journal of Food Science*, 66:754 – 757, 2001.
- [28] G. O. Sigge, C. F. Hansmann and E. Joubert, Effect of storage conditions, packaging material and metabisulphite treatment on the colour of dehydrated green bell peppers (*Capsicum annum* L.). *Journal of Food Quality*, 24:205 – 218, 2001.
- [29] A. Silvica, G. Sansheng, P. M. Brezeanu, C. Brezeanu, and M. Calin, Studies on culture behavior of some sweet pepper varieties (*Capsicum annum* L). *Journal of Horticulture, Forestry and Biotechnology*, 15:197-200, (2011).